## MODERN CLASSIFICATION OF CUTANEOUS PSEUDOLYMPHOMAS

S. Shtilionova<sup>1</sup>, P.Drumeva<sup>1</sup>, M. Balabanova<sup>2</sup>, I. Krasnaliev<sup>3</sup>

- 1) Department of dermatology and venereology, Medical University Varna
- 2) Department of dermatology and venereology, Medical University Sofia
- 3) Department of pathology, Medical University Varna

#### SUMMARY

Systematizing of different diseases and skin disorders is quite difficult. Different authors make classifications according to numerous criteria. Pseudolymphomas are reactive lymphocytic proliferations that appear in the skin and resemble a malignant lymphoma.

Different classifications exist for cutaneous pseudolymphomas. The broadest classification differentiates them into 6 groups, but the most commonly used is the classification based on clinical and morphological criteria.

According to the latest classification (1998), they are divided into cutaneous T-cell pseudolymphomas and - cutaneous B-cell pseudolymphomas.

The evidence from the histopathological analysis is of utmost important for the accurate diagnosis.

Key words: Cutaneous pseudolymphomas, Classification

Cutaneous pseudolymphomas are benign lymphoproliferating processes resembling malignant lymphomas clinically and histologically, in particular (1, 2). They present an accumulation of lymphocytes in response to various stimuli (infections, drugs, insect bites, etc.). This is a heterogeneous group of dermatoses with clinical manifestations varying from tumor-like nodes to flat cell infiltrates. Many authors divide them into pseudolymphomas in the narrow sense and pseudolymphomas in the broad sense of the term (4, 5). According to the type of the cell infiltrates, the cutaneous lymphomas are divided into T-cell, B-cell and mixed (3).

Various classifications exist for cutaneous pseudolymphomas (1, 2, 3, 4). One of the relatively clearest and most frequently used in practice is Rijlaarsdam& Willemze's classification based on clinical and morphological criteria.

### 1. Cutaneous T-cell pseudolymphomas

a) Primarily with stripe-like infiltration (the majority of cases)

Lymphomatoid drug eruption (most cases); Lymphomatoid contact dermatitis; Actinic reticuloid; Nodular scabies (individual cases); Idiopathic forms;

Clonal cutaneous T-cell pseudolymphomas.

b) Primarily with nodular infiltration (a small percentage of the cases)

Drug-induced – mainly by anti-convulsive drugs (individual cases);

Persistent nodules after insect bites; Nodular scabies (the majority of cases).

# 2. Cutaneous B-cell pseudolymphomas (with nodular infiltration)

Cutaneous lymphocytoma from Borrelia burgdorferi; Cutaneous lymphocytoma after antigens injection; Cutaneous lymphocytoma resulting from tattoo; Cutaneous lymphocytoma after Herpes zoster; Idiopathic forms; Clonal cutaneous B-cell pseudolymphomas.

Another, also very common classification, is the one dividing them into 6 groups (Burg&Braun-Falco; Kerl&Smole) (5, 6):

### CLASSIFICATION

*1. Infiltrations from non-lymphoid cells* – tumor cells in neuroblastoma and in the tumor of Merkel's cells can resemble a malignant lymphoma.

2. Neoplasm rich in lymphocytes – for instance, cutaneous lymphadenoma (variant of trichoblastoma) can be so rich in lymphocytes that its epithelial component can be very difficult to identify.

3. Stroma reaction in epithelial displasia and malignant neoplasms of the soft tissues – lymphoid variants of actinic keratoses are well-known; neoplasms such as basal cell carcinoma and dermatofibroma can cause extremely powerful stroma reactions, in which the primary neoplasm can be difficult to detect.

4. Diseases which are not directly related to the skin – they comprise a long list and include Rosai-Dorfmann's

disease, Castleman's disease, Kikuchi's disease, inflammatory pseudotumor and malacoplacia.

5. Classical dermatological diseases resembling cutaneous lymphoma – again the list is rather long, most frequently related to autoimmune diseases of the connective tissue. This group includes lychan sclerosus et atrificus, variants of lupus erythematosus, atypical lymphocyte lobular paniculitis, lymphomatoid dermatitis and lymphomatoid folliculitis.

**6.** Specific cutaneous pseudolymphoma units – acral pseudolymphoma angiokeratoma in children (APACHE), palpable arciform migratory erythema, angiolymphoid hyperplasia with eosiniphilia and Kimura's disease. The

lymphomatoid papulosis is not classified under this group any more but falls into the group of the primary cutaneous lymphomas.

### **DISCUSSION:**

The concept of cutaneous lymphomas provides the basis for the understanding and the identification of the benign and malignant cutaneous lymphoid infiltrations. Even following accurate histological investigations, many of the lymphoid infiltration processes remain unclearly defined as benign or malignant. The diagnosis of the cutaneous lymphoid infiltrations requires series of biopsies and examinations.

### **REFERENCES:**

1. Crowson AN. Cutaneous pseudo-lymphoma: a review. Fitzpatrick's J Clin Dermatol 1995; 3: 43-55.

2. Brodell RT, Santa Cruz DJ. Cutaneous pseudolymphomas. Dermatol Clin. 1985 Oct;3(4):719-734. http:// www.ncbi.nlm.nih.gov/pubmed/3916177

3. Dorfman RF, Warnke R. Lympha-denopathy simulating the malignant lymphomas. Hum Pathol 1974 Sep;5(5):519-550. http:// www.ncbi.nlm.nih.gov/pubmed/4137045

4. Gartman H. Generalisierte pseudo-lymphoma der haut. Hautarzt 1986; 55: 166.

5. Medeiros LJ, Picker LJ, Abel EA, et al. Cutaneous lymphoid hyperplasia. Immunologic characteristics and assessment of criteria recently proposed as diagnostic of malignant lymphoma. J Am Acad Dermatol 1989 Nov;21(5 Pt 1):929-942. http://www.ncbi.nlm.nih.gov/pubmed/ 2808829

6. Ploysangam T, Breneman DL, Mutasim DF. Cutaneous pseudolymphomas. J Am Acad Dermatol 1998; 38: 877-895; quiz 896-897.

7. Burg G, Braun-Falco O. Cutaneous pseudolymphomas. In: Cutaneous lymphomas. Berlin: Springer -Verlag, 1983, 415-464.

8. Burg G, Kerl H, Schmoeckel C. Differentiation between malignant B-cell lymphomas and pseudolymphomas of the skin. J Dermatol Surg Oncol. 1984 Apr; 10(4):271-275. http:// www.ncbi.nlm.nih.gov/pubmed/6608539

9. Kerl H, Smolle J. Classification of cutaneous pseudolymphomas. Curr Probl Dermatol 1990; 19: 167-175.

10. Rijlaarsdam JU, Bruynzeel DP, Vos W, Meijer CJ, Willemze R. Immunohistochemical studies of lymphadenosis benigna cutis occurring in a tattoo. Am J Dermatopathol. 1988 D e c ; 1 0 ( 6 ) : 5 1 8 - 5 2 3 . h t t p : // www.ncbi.nlm.nih.gov/pubmed/3218720

11. Rijlaarsdam U, Scheffer E, Meijer CJ, Kruyswijk MR, Willemze R. Mycosis fungoides-like lesions associated with phenytoin and carbamazepine therapy. J Am Acad Dermatol 1991 Feb;24(2 Pt 1):216-220. http://www.ncbi.nlm.nih.gov/ pubmed/1826111

12. Rijlaarsdam JU, Willemze R. Cutaneous pseudolymphomas: classifica-tion and differential diagnosis. Semin Dermatol 1994 Sep;13(3):187-196. http://www.ncbi.nlm.nih.gov/pubmed/ 7986687